

Kondo effect in a double quantum-dot molecule under the effect of an electric and magnetic field

Gustavo. A. Lara,¹ Pedro A. Orellana,² and Enrique V. Anda³

¹*Departamento de Física, Universidad de Antofagasta, Casilla 170, Antofagasta, Chile.*

²*Departamento de Física, Universidad Católica del Norte, Casilla 1280, Antofagasta, Chile.*

³*Departamento de Física, P. U. Católica do Rio de Janeiro, C.P. 38071-970, Rio de Janeiro, RJ, Brazil.*

Electron tunneling through a double quantum dot molecule, in the Kondo regime, under the effect of a magnetic field and an applied voltage, is studied. This system possesses a complex response to the applied fields characterized by a tristable solution for the conductance. The different nature of the solutions are studied in and out thermodynamical equilibrium. It is shown that the interdot coupling and the fields can be used to control the region of multistability. The mean-field slave-boson formalism is used to obtain the solution of the problem.

PACS numbers: PACS number(s): 73.21.La; 73.63.Kv; 85.35.Be

The electron tunneling through a double-quantum-dot (DQD) molecule in the Kondo regime has received much attention in the last years.^{1,2,3,4,5,6,7,8,9} Recently Jeong et al.¹ reported the observation of a coherent Kondo effect in a double quantum dot, which is a direct experimental realization of a two-impurity Anderson model.^{2,3,4,5,6,7,8} The strength of the coupling between the QD's determines the character of the molecular electronic states and the transport properties of the artificial molecule. In the tunneling regime, the states are extended across the entire system and can be constructed from a coherent state based on the bonding or anti-bonding levels of the quantum dots. These many-body molecular states are observed in a transport experiment as a differential conductance of two peaks, a consequence of the splitting of the Kondo resonance.

Recently Aono and Eto studied theoretically the problem of a DQD in thermodynamical equilibrium, under the effect of a magnetic field oriented parallel to the current direction.⁶ They calculated the magnetoconductance and found a sudden drop of the conductance as a function of the field, when $t_c > \Gamma$, where t_c is the interdot interaction and Γ the resonant width of the dot levels due to coupling among them and the contacts. Some aspects of the non-equilibrium transport properties of a DQD in the Kondo regime, under the effect of an applied voltage, has recently been studied⁹. When the inter-dot coupling is greater than the level broadening and there is an external voltage applied to the system, the conductance shows a tristable behavior.

In this work we analyze the physics derived from the complex behavior that characterizes the electron tunneling through a DQD molecule in the Kondo regime under the simultaneous effect of a magnetic and an electric field. We show that the discontinuity of the conductance found

by Aono and Eto,⁶ increasing the magnetic field when the system is in equilibrium, is a result of a tristable behavior of the magnetoconductance. This results is similar to the behavior the system has when it is out of equilibrium, due to the application of an external voltage.⁹

We consider a DQD connected in series attached to two leads. The physics associated to this artificial molecule electrically connected can be readily understood in terms of two impurities Anderson Hamiltonian where the impurities located in the site 0 and 1 are the quantum dots. The leads are represented by a one dimensional tight-binding Hamiltonian connecting the dots to two particle reservoirs characterized by Fermi levels μ_L and μ_R respectively. The left (right) dot is connected to the left (right) lead by the hopping $V_{L(R)}$ and the dots are connected among themselves by the interdot tunneling coupling t_c . Each dot α has a single energy ε_α ($\alpha = 0, 1$). Under an applied magnetic field the energy levels of the dots are split by the Zeeman effect into $\varepsilon_\alpha \pm \sigma g \mu_B B$. We adopt the two-fold degenerate Anderson Hamiltonian in the limit $U \rightarrow \infty$, that is diagonalized using the mean-field slave-boson formalism.^{10,11,12} The double occupancy in each dot is forbidden and the inter-dot Coulomb interaction neglected. We introduce the slave-boson operator $b_\alpha^\dagger$ that creates an empty state and a fermion operator $f_{\alpha\sigma}$ that annihilates a single occupied state with spin σ . To eliminate the possibility of double occupancy we impose the constraint $Q_\alpha \equiv \sum_\sigma f_{\alpha\sigma}^\dagger f_{\alpha\sigma} + b_\alpha^\dagger b_\alpha = 1$. The annihilation operator of an electron in the dot α is $c_{\alpha\sigma} = b_\alpha^\dagger f_{\alpha\sigma}$. In the mean field approximation the bosonic operators $b_\alpha^\dagger$ and b_α are replaced by their expectation values, $\langle b_\alpha \rangle = \tilde{b}_\alpha \sqrt{2} = \langle b_\alpha^\dagger \rangle = \tilde{b}_\alpha^\dagger \sqrt{2}$. Hence the Hamiltonian of the DQD molecule connected to leads plus constraints is written as,

$$\begin{aligned}
H = & H_{lead} + \sum_{\alpha=0,1,\sigma} (\tilde{\varepsilon}_\alpha + \sigma g \mu_B B) f_{\alpha\sigma}^\dagger f_{\alpha\sigma} + \tilde{V}_L \sum_{\sigma} (c_{-1\sigma}^\dagger f_{0\sigma} + f_{0\sigma}^\dagger c_{-1\sigma}) + \tilde{V}_R \sum_{\sigma} (f_{1\sigma}^\dagger c_{2\sigma} + c_{2\sigma}^\dagger f_{1\sigma}) \\
& + \tilde{t}_c \sum_{\sigma} (f_{0\sigma}^\dagger f_{1\sigma} + f_{1\sigma}^\dagger f_{0\sigma}) + \sum_{\alpha=0,1} \lambda_\alpha (\tilde{b}_\alpha^\dagger \tilde{b}_\alpha - 1).
\end{aligned} \tag{1}$$

where $\tilde{\varepsilon}_\alpha = \varepsilon_0 + \lambda_\alpha$, $\tilde{V}_L = V_L \tilde{b}_0$, $\tilde{V}_R = V_R \tilde{b}_1$, $\tilde{t}_c = t_c \tilde{b}_0 \tilde{b}_1$, the λ_α are Lagrangian multipliers, which guarantee the constraint conditions on the Q_α .

The first term on the right-hand side of Eq.1 represents the electrons in the left and right leads;

$$H_{lead} = \sum_{\sigma, i \neq 0,1} \varepsilon_i n_{i\sigma} + t \sum_{\langle ij \neq 0,1 \rangle \sigma} c_{i\sigma}^\dagger c_{j\sigma}. \tag{2}$$

where the operator $c_{i\sigma}^\dagger$ creates an electron in the site i with spin σ , ε_i is the site energy and t is first-neighbor hopping in the leads. Here H_{lead} is a free-particle Hamiltonian with eigenfunctions of the Bloch type,

$$|k, \sigma\rangle = \sum_j e^{ikja} |j, \sigma\rangle, \tag{3}$$

where $|k, \sigma\rangle$ is the momentum eigenstate with spin σ and $|j\rangle$ is a Wannier state localized at site j of spin σ . The dispersion relation associated with these Bloch states reads

$$\varepsilon = 2t \cos(kd). \tag{4}$$

The model has an energy band, extending from $-2t$ to $+2t$, with the first Brillouin zone expanding the k interval $[-\pi/d, \pi/d]$.

The stationary state of the complete Hamiltonian H with energy ε_k can be written as

$$|\psi_{k\sigma}\rangle = \sum_j a_{j\sigma}^k |j, \sigma\rangle, \tag{5}$$

where the coefficients $a_{i\sigma}^k$ are given by,

$$a_{j\sigma}^k = \langle j, \sigma | \psi_{k\sigma} \rangle. \tag{6}$$

We obtain the following eigenvalue equations for the Wannier amplitudes $a_{j\sigma}^k$

$$\begin{aligned}
\varepsilon a_{j,\sigma}^k &= \langle j, \sigma | H | \psi_{k\sigma} \rangle \\
\varepsilon a_{j,\sigma}^k &= \varepsilon_j a_{j,\sigma}^k + t(a_{j-1,\sigma}^k + a_{j+1,\sigma}^k) \quad (j \neq -1, 0, 1, 2), \\
\varepsilon a_{-1(2),\sigma}^k &= \varepsilon_{-1(2)} a_{-2(2),\sigma}^k + \tilde{V}_{L(R)} a_{0(1),\sigma}^k + t a_{-2(3),\sigma}^k, \\
\varepsilon a_{0(1),\sigma}^k &= (\tilde{\varepsilon}_{0(1)} + \sigma g \mu_B B) a_{0(1),\sigma}^k + \tilde{V}_{L(R)} a_{-1(2),\sigma}^k \\
&\quad + \tilde{t}_c a_{1(0),\sigma}^k,
\end{aligned} \tag{7}$$

where $a_{j,\sigma}^k$ is the probability amplitude to find the electron in site j and state k with spin σ .

The four parameters ($\tilde{b}_0$, $\tilde{b}_1$, λ_0 , λ_1) are determined minimizing the expectation value of the Hamiltonian (1). They satisfy the set of equations:

$$\begin{aligned}
\tilde{b}_{0(1)}^2 + \frac{1}{2} \sum_{k,\sigma} |a_{0(1),\sigma}^k|^2 &= \frac{1}{2}, \\
\frac{\tilde{V}_{L(R)}}{2} \sum_{k,\sigma} \text{Re}(a_{-1(2),\sigma}^{k*} a_{0(1),\sigma}^k) + \frac{\tilde{t}_c}{2} \sum_{k,\sigma} \text{Re}(a_{1(0),\sigma}^{k*} a_{0(1),\sigma}^k) + \tilde{\lambda}_{0(1)} \tilde{b}_{L(R)}^2 &= 0.
\end{aligned} \tag{8}$$

The sum over the wave vector k cover all the occupied states. The resulting equations are nonlinear due to the renormalization of the localized levels in the dots, the interdot coupling tunneling and the coupling tunneling between the quantum dots and the leads.

In order to study the solutions of equations (7) and (8) we assume a plane wave incident from one lead with an

amplitude I and with a partial reflection amplitude R . The waveform at the other lead is a simple plane wave with intensity given by the transmission amplitude T . Taking this to be the solution of the system at infinite,

for positive k , we can write

$$\begin{aligned} a_{j\sigma}^k &= I_\sigma^+ e^{ikdj} + R_\sigma^+ e^{-ikdj}, & j < 0 \\ a_{j\sigma}^k &= T_\sigma^+ e^{ikRdj}, & j > 1 \end{aligned} \quad (9)$$

and those with negative k

$$\begin{aligned} a_{j\sigma}^{-k} &= I_\sigma^- e^{-ikdj} + R_\sigma^- e^{ikdj}, & j > 1, \\ a_{j\sigma}^{-k} &= T_\sigma^- e^{-ikLdj}, & j < 0. \end{aligned} \quad (10)$$

These functions describe free electrons approaching the scattering potential from $j = -\infty$ and $j = +\infty$, respectively, with transmission and reflections amplitudes T_σ^+ and R_σ^+ for $k > 0$ and T_σ^- and R_σ^- for $k < 0$.

The solution of equations (7) can be obtained through an adequate iteration from right to left for $k > 0$, and from left to right for $k < 0$. For a given transmitted amplitude, the associated reflected and incident amplitudes may be determined by matching the iterated function to the proper plane wave at the left lead for $k > 0$ and at the right lead for $k < 0$. The transmission amplitude $t_\sigma = T_\sigma^+/I_\sigma^+$ and $\tilde{t}_\sigma = T_\sigma^-/I_\sigma^-$ are obtained using an iterative procedure. The total transmission probability with a spin σ and energy $\varepsilon = 2t \cos(kd)$ given by $T(\varepsilon) = \sum_{\sigma=\uparrow,\downarrow} |t_\sigma(\varepsilon)|^2$. Then the linear conductance at zero temperature is written as,

$$G = \frac{2e^2}{h} T. \quad (11)$$

The equations (7) and (8) are nonlinear and require a self-consistent solution that is obtained using a conjugate gradient algorithm. In this work, we are interested in analyzing all the stationary solutions for the conductance. As $G(B)$ and $B(G)$ are multivalued functions, the multiple solutions for the quantities $\tilde{b}_0, \tilde{b}_1, \lambda_0, \lambda_1$ are obtained by fixing the magnetic field B and obtaining the value of the conductance G that corresponds to the solution of the equations of motion of the system. This method permits us to obtain the complete $G - B$ curve.

We study first a model which consists of two leads equally connected ($V_L = V_R = V_0$) to both sides of a DQD dots in equilibrium with chemical potential $\mu_L = \mu_R = 0$, $t = 30 \Gamma_0$, $V_0 = 5.48 \Gamma_0$, $\varepsilon_0 = \varepsilon_1 = -3.5 \Gamma_0$ (Kondo regime with $T_K \approx 10^{-3} \Gamma_0$ with $\Gamma_0 = 2\pi V_0^2 \rho(0)$).

Fig 1 shows the linear conductance versus the applied magnetic field for different values of t_c with $t_c \leq \Gamma_0$. For small values of t_c , $t_c \ll \Gamma_0$ our results agrees with those obtained by Aono and Eto. The magnetoconductance is negative in this region. However for greater values of t_c , $t_c \sim \Gamma_0$ another behavior of the magnetoconductance starts. Within a range of values of the magnetic field there is a tristability behavior with three stationary solutions of the conductance for each value of the magnetic field. Fig 2 shows the results for $t_c > \Gamma_0$. As in the results found by Aono and Eto, at a critical value $B_{c\uparrow}$, there




FIG. 1: G - B curve for a) $t_c = 0.2\Gamma_0$, b) $t_c = 0.6\Gamma_0$, c) $t_c = 0.8\Gamma_0$ and d) $t_c = \Gamma_0$

would be a discontinuous reduction of the conductance when B is increased and other $B_{c\downarrow}$ ($B_{c\uparrow} > B_{c\downarrow}$) where the conductance would rise discontinuously, when B is reduced. Using this procedure, the interior of the tristability remains inaccessible to the experiment. However, the internal shape could eventually be reached, as it is for the case of double barrier heterostructures, if it were possible to employ a load line in the experimental measurement to go along all the points of the $G - B$ curve. From this point of view the electrons goes along a non-linear potential, that is magnetic field dependent. In this case the magnetic field plays the role that the external electric potential plays in the case of a standard double barrier heterostructure, where the current has as well a multistable behavior.¹³




FIG. 2: Linear conductance G versus the Zeeman splitting $g\mu_B B$ for a) $t_c = 1.0\Gamma_0$, b) $t_c = 1.2\Gamma_0$, c) $t_c = 1.6\Gamma_0$ and d) $t_c = 1.8\Gamma_0$

Increasing the magnetic field the tristability probably arises from a many-body ground state that changes its character due to a energy crossing. It is a singlet of the Kondo type when there is no applied magnetic field, while for a critical field onwards the ground state results to be

the component of the triplet wave function, (two parallel spins one at each dot) with spin projection parallel to the magnetic field. In the parameter space this two ground states have their own basin of attraction constituting metastable states when they loose the condition of been the ground state of the system. As soon as the metastability is lost increasing or decreasing the field, the system abruptly relaxes to the ground state producing a discontinuous behavior of the current.¹⁹

It is very interesting to study these non-linear effect when the system is under the simultaneous action of an external magnetic and electric field. In this case the DQD is out of equilibrium as it operates within two chemical potentials $\mu_L = V/2$ and $\mu_R = -V/2$ defined by the applied dc bias voltage V . Imposing a fixed value of the magnetic field, once the amplitudes $a_{j,\sigma}^k$ are known, the current can be numerically obtained from,

$$J = \frac{2e}{\hbar} \tilde{V}_L \sum_{k,\sigma} \text{Im}\{a_{-1,\sigma}^{k*} a_{0,\sigma}^k\}. \quad (12)$$




FIG. 3: J - V characteristic curve for $t_c = \Gamma_0$ for various values of the Zeeman splitting $g\mu_B B$. a) $g\mu_B B = 0.0$, b) $g\mu_B B = 0.40T_K$, c) $g\mu_B B = 0.46T_K$ and d) $g\mu_B B = 0.47T_K$

We study first the problem when $t_c = \Gamma$. For this value of t_c , in the absence of an external magnetic field, the system is at the threshold of having a manifestation of the non-linearity produced by correlation. For a regime in which $g\mu_B B \ll T_K$, figure 3 shows that the magnetic field reduces the height of the peak and valley of the $J - V$ characteristic curve. However, for larger magnetic fields, $g\mu_B B < T_K$, a tristability behavior of the current is promoted. For greater values of t_c ($t_c = 1.4\Gamma$), where the system has already a highly non-linear behavior in the absence of magnetic field, the shape of the tristability is highly dependent on B . As shown in figure 4, here as well, a remarkable increase of the peak to valley ratio is obtained in the regime $g\mu_B B < T_K$. This ratio goes from about 20 at zero magnetic field to about 2000 at $g\mu_B B = 0.333T_K$. These phenomena are a result of the reduction and eventual elimination of the

Kondo correlation when the magnetic field is of the order of the Kondo temperature. The Zeeman effect destroys the coherent behavior of the two QD's produced by the molecular Kondo coupling, the electrons localize at each dot occupying levels that are mismatched by the applied potential and consequently the current goes to zero.




FIG. 4: J - V characteristic curve for $t_c = 1.4\Gamma_0$ for various values of the magnetic field. a) $g\mu_B B = 0$, b) $g\mu_B B = 0.2T_K$, c) $g\mu_B B = 0.3T_K$ and d) $g\mu_B B = 0.33T_K$

In non-equilibrium the nonlinear behavior described above has similarities with the well-known bistability present in three dimensional resonant tunneling double barrier systems and multistabilities in doped superlattices.^{9,14,15,16,17,18} In three dimensional tunneling double barrier systems, the Coulomb interaction pins the renormalized energy level at the well when the applied voltage V is augmented maintaining the system at resonance. However, when V goes above a critical value, the abrupt leakage of the charge accumulated in the well takes the system out of resonance. For doped superlattices, at large carrier densities, the internal potential profile produced by the applied voltage breaks up into domains giving rise to a multistable behavior in the current.^{15,16,17} In both systems, the non-linear deformation of the potential profile due to the electron-electron interaction is responsible for the phenomenon. Although in our case the Kondo effect is a strong many body effect of a different nature in comparison with the previous systems, the multistability is as well a consequence of the non-linear potential profile deformations that the electrons suffer when the external potential is modified.

In conclusion, we have investigated the magnetoconductance in a DQD molecule in the Kondo regime. We have shown that the magnetoconductance has a tristable behavior depending of the ratio of t_c/Γ and the value of the magnetic field that controls the non-linearity of the system. This evidence was obtained using a mean-field approach that neglects fluctuations, appropriate for large spins as it is the first term of an $1/N$ expansion, where N is the spin dimension. Some experiments have shown²¹ that it is possible to design dots with a configuration that, due to the exchange interaction, the parallel

spin coupling following Hund's rule gives rise to a total dot spin $S_t > 1/2$. For this cases our results are more reliable as fluctuations are less important. Although we believe that our results are robust against them, a study that goes beyond mean field is necessary to confirm our

conclusions.

Work supported by grants Milenio ICM P99-135-F, FONDECYT Grants 1020269 and 7020269 and Red IX.E "Nanoestructuras para la Micro y Optoelectronica Sub-Programa IX "Microelectronica" Programa CYTED.

-
- ¹ H. Jeong, A.M. Chang and M.R. Melloch, Science **293** 2221 (2001).
 - ² R.Aguado and D.C. Langreth Phys.Rev.Lett. **85**,1946 (2000).
 - ³ Antoine Georges and Yigal Meir, Phys. Rev. Lett. **82**, 3508 (1999).
 - ⁴ C.A. Busser, E.V. Anda, A.L. Lima, M. A.Davidovich and G. Chiappe, Phys. Rev. B **62**, 9907 (2000).
 - ⁵ Tomosuke Aono and Mikio Eto, Phys. Rev. B **63**,125327 (2001).
 - ⁶ Tomosuke Aono and Mikio Eto, Phys. Rev. B **64**,073307 (2001).
 - ⁷ Wataru Izumida and Osamu Sakai, Phys. Rev. B **62** 10260 (2000).
 - ⁸ T. Ivanov Europhys. Lett. **40**, 183 (1997).
 - ⁹ Pedro A. Orellana, G.A. Lara, Enrique V. Anda, Phys. Rev.B **65** 155317 (2002),Gustavo A. Lara, Pedro A. Orellana, Enrique V. Anda, Sol. State Comm. **125** 165 (2003).
 - ¹⁰ S.E. Barnes, J. Phys. F **6**, 1375 (1976); **7**, 2637 (1977).
 - ¹¹ P. Coleman, Phys. Rev. B **29**, 3035 (1984); **35**, 5072 (1987).
 - ¹² N. Read and D.M. Newns, J. Phys. C, 3273 (1983); Adv. Phys. **36**, 799 (1988).
 - ¹³ A.D Martin, M.K.F. Lerch, P.E. Simmonds and E. Eaves, Appl. Phys. Lett. 64, 1248 (1994).
 - ¹⁴ V.J. Goldman, D.C.Tsui and J.E. Cunningham, Phys. Rev. Lett. **58**, 1256 (1987).
 - ¹⁵ F. Prengel, A. Wacker, and E. Scholl, Phys. Rev. B **50**, 1705 (1994).
 - ¹⁶ J. Kastrup,R. Hey, K. H. Ploog, H. T. Grahn, L. L. Bonilla, M. Kindelan, M. Moscoso, A. Wacker, and J. Galán, Phys. Rev. B **55**, 2476 (1997).
 - ¹⁷ R. Aguado,G. Platero, M. Moscoso, and L. L. Bonilla, Phys. Rev. B **55**, R16053 (1997).
 - ¹⁸ P.L. Pernas, F. Flores and E.V. Anda, Phys. Rev. B **47**, 4779 (1993).
 - ¹⁹ E. V. Anda ,V. Ferrari and G. Chiappe, J.Phys.:Condens.Matter **9**, 1095 (1997).
 - ²⁰ B.A. Jones, B.G. Kotliar, A. J. Millis, Phys. Rev. B **39**, R3415 (1989).
 - ²¹ S.Sasaki, S. De Franceschi, J.M. Elzerman, W.G. van de Wiel, M. Eto, S. Tarucha and L. P. Kowenhoven, Nature **405**, 764 (2000).